

J. Wolff

STICHTING ONDERZOEK WERELDVOEDSELVOORZIENING

CENTRE FOR WORLD FOOD STUDIES

THE CONCENTRATION OF MACRO-NUTRIENTS IN (SUB)TROPICAL CROPS
OF MINOR IMPORTANCE

by

K. NIJHOF

- Staff working paper SOW-87-17 -

Centre for World Food Studies
c/o CABO
P.O. Box 14, 6700 AA Wageningen
The Netherlands

November 1987

Staff working papers are informal notes on research, conducted by the
Centre For World Food Studies. They should not be quoted without the
prior permission of the Centre.

INTRODUCTION

In order to predict crop yields in situations that crop growth is limited by nutrient availability, a system for quantitative evaluation of the fertility of tropical soils (QUEFTS) has been developed by Janssen et al (1986 and 1987).

The inputs required for the application of the QUEFTS system are, apart from chemical soil data, the minimum and maximum concentrations of the macronutrients N, P and K in the economic product of a crop and in its residues and the dry matter distribution. Initially, these data were established for maize only, as QUEFTS was originally developed for this crop. However, QUEFTS can be adapted to any crop if the ranges in N, P and K concentrations in residues and economic products and the Harvest Index are known. Such data have recently been collected by the Centre for World Food Studies for the major (sub)tropical crops (Nijhof, 1987). Data for crops of minor importance are still lacking however.

In this study data are collected on the ranges in N, P and K concentrations as well as Harvest Indices for the "minor" (sub)tropical crops, particularly fruit and vegetable crops. These data are used for the calculation of yield-nutrient uptake relations.

METHODS

A large number of articles with data on the N, P and K concentrations and the dry matter distribution and concentration in different crops is reviewed (Appendices 1 and 2). These data are used to establish the minimum and maximum nutrient concentrations. They are derived for mature (i.e. to be harvested), complete plants (excluding roots) and the corresponding concentrations in economic products and residues are determined afterwards (see Table 1.) Extreme high or low concentrations are often found in one part of the crop, but are only used as minimum or maximum concentrations if they are accompanied by high or low concentrations in the other crop

parts. The established ranges cover 90% of the values found, i.e. the upper and lower extremes are excluded. Correlations between the N and P concentrations in a plant part as well as between the concentrations in economic product and residue are computed for all crops. Both types of correlations, if they proved to be strong, are used to determine the range of concentrations of the least documented nutrient and in the least documented plant part respectively.

Harvest Indices are calculated as the means of the values found. These indices are used to establish yield-nutrient uptake relations at minimum and maximum nutrient concentrations.

The Harvest Indices and nutrient ranges of pineapple and banana are determined for the mother plants only, i.e. ignoring the daughter plants and/or young shoots. The data collected for castor beans refer to annual varieties, although this crop can be grown as perennial too.

RESULTS

The values for the minimum and maximum N, P and K concentrations in the crops and their mean Harvest Indices are presented in the Tables 1 and 2.

Information on the nutrient concentrations in the residues of chillies and castor is rare. These values are estimated using data of the residues of other Capsicum species and cassava tops, respectively.

Reliable yield-nutrient uptake relations should include the

Table 1. Nutrient concentration in economic products and crop residues of (sub)tropical crops, as fraction of N, P and K on dry matter basis.

crop organ	N min-max %	P min-max %	K min-max %
Banana/Plantain			
bunch	0.60-1.50	0.08-0.18	2.10-4.00
fruits	0.60-1.50	0.08-0.18	2.00-3.40
residues	0.70-1.40	0.07-0.17	1.50-8.60
Cabbage (white cabbage)			
heads + core's	1.80-4.20	0.19-0.70	1.50-4.50
residue	1.40-2.50	0.14-0.39	0.80-3.70
Carrot			
roots	0.50-2.30	0.10-0.60	0.70-4.30
tops	0.70-3.60	0.10-0.60	0.80-6.00
Castor (dwarf varieties)			
seeds	2.90-4.10	0.30-0.80	0.60-2.50
shell approximation	0.90-0.90	0.22-0.22	0.90-0.90
haulms *	0.50-1.80	0.08-0.50	0.45-1.80
straw	0.50-1.70	0.09-0.50	0.50-1.70
Chillies			
pods (red ripe)	1.20-3.50	0.25-0.50	1.40-3.90
straw	1.00-3.50	0.11-0.40	1.20-4.40
Garlic			
bulbs	0.90-2.60	0.17-0.50	0.70-1.30
tops	1.40-3.50	0.16-0.32	1.40-4.70
Okra			
pods	1.50-3.40	0.20-0.70	0.80-4.30
straw	0.50-2.40	0.07-0.26	0.90-3.60
Spinach: whole plants **			
Amaranthus spp.	2.50-5.50	0.60-0.80	3.70-7.80
flowering Amaranth	0.70-3.90	0.19-0.60	3.60-5.40
Indian Spinach ***	2.00-4.70	0.24-0.35	2.90-3.30
Kang Kong	3.20-6.30	0.23-0.60	3.10-6.60

* haulms: stem + leaves

straw: stem + leaves + shells

** Young plants if not otherwise indicated

*** Basella Alba

crop organ	N	P	K
	min-max %	min-max %	min-max %
Pigeon Pea			
seeds	2.80-4.30	0.20-0.50	0.70-1.20
shells	0.70-1.80	0.04-0.12	0.30-1.00
haulms *	0.70-2.00	0.07-0.20	0.30-1.10
straw	0.70-1.90	0.06-0.15	0.30-1.10
Pineapple			
fruits	0.45-2.00	0.04-0.18	0.70-2.70
fruit/crown/stalk	0.50-2.70	0.04-0.18	0.60-2.90
residue	0.70-3.60	0.07-0.16	0.60-3.70
Tomatoes			
fruits	1.70-3.50	0.15-0.90	1.50-6.00
straw	1.10-3.30	0.12-0.80	1.80-5.50

* haulms: stem + leaves

straw: stem + leaves + shells

** Young plants if not otherwise indicated

*** Basella Alba

nutrients taken up in the economic products and residues as well as the nutrients lost by leaf shedding. Therefore, Harvest Indices should also include the fallen leaves. Data on the amount of shedded leaves are lacking in the literature for a few crops and the Harvest Index has to be estimated from the data collected for the Harvest Index excluding the shedded leaves. This is done by taking 75-85 percent of the means of the collected Harvest Indices, depending on the stem/leaf ratio of the crop and its maturity at harvesting stage.

Table 3 presents the calculated yield-nutrient uptake relations. The data of Table 3 are converted into yields of fresh marketable products as presented in Table 4.

REFERENCES

Janssen, B.H., F.C.T. Guiking, D. van der Eijk, E.M.A. Smaling and H. van Reuler (1986) A new approach to evaluate the chemical fertility of tropical soils. Proceedings ISSS Congress, Hamburg, Volume 3: 791-792.

Janssen, B.H., F.C.T. Guiking, D. van der Eijk, E.M.A. Smaling, J. Wolf and H. van Reuler (in prep.) A system for quantitative evaluation of the fertility of tropical soils (QUEFTS).
 Nijhof, K. (1987) The concentration of macro-elements in economic products and residues of (sub)tropical field crops. SOW Staff Working Paper 87-07.

Table 2. Harvest Indices (HI) of (sub)tropical crops, calculated for oven-dry crop organs and the dry matter (DM) concentration of their economic products.

crop economic product	HI excl. shed leaf	HI incl. shed leaf	DM % of econo- nomic product.
	%	%	%
Banana/Plantain			
bunch	39	35	-
fruit	37	34	21
Cabbage (white cabbage)			
heads + core's	47	45	7
Carrot			
roots	70	51	10
Castor (dwarf varieties): approximations			
pods	29	39 *	-
seeds	19	15 *	86
oil	9	7 *	100
Chillies			
pods (red ripe)	48	36 *	87
Garlic			
bulbs	50	40 *	37
Okra			
pods	26	22	10
Pigeon Peas			
pods	28	22	-
seeds	17	14	84
Pineapple			
fruits	25	23	11
fruit/stalk/crown	36	33	-
Spinach: edible leaves + stems **			
Amaranthus spp	69	69	10
flowering Amaranth	21	17 *	17
Indian Spinach ***	100	100	7
Kang Kong	100	100	9
Tomato			
fruits	63	53	6

* estimation

** Young plants unless otherwise indicated.

*** Basella Alba

Table 3. Economic product-nutrient uptake relations of (sub)tropical crops at minimum and maximum N, P and K concentration (kg dry matter per kg nutrient uptake).

crop	N		P		K	
	max-min		max-min		max-min	
Banana/Plantains						
bunches	25	-53	204-481		5.1-21	
fruits	24	-52	198-467		4.9-20	
Cabbage (white cabbage)						
heads + core's	14	-28	85-227		11	-40
Carrot						
roots	17	-85	85-510		13	-68
Castor (dwarf varieties)						
pods	12	-27	43-195		13	-46
seeds	7.3-17		28-124		8.2-28	
Chillies						
pods (red ripe)	10	-34	83-224		8.5-28	
Garlic						
bulbs	13	-33	103-326		12	-35
Okra						
pods	8.4-31		62-223		5.9-25	
Pigeon Pea						
pods	10	-23	87-282		20	-63
seeds	6.3-14		54-175		13	-39
Pineapple						
fruit/stalk/crown	9.1-46		179-466		8.5-47	
fruits	7.1-36		140-365		6.6-37	
Spinach: edible leaves + stems						
Amaranth young	13	-29	86-115		8.9-19	
flowering Amaranth	4.4-24		28- 90		3.2- 4.7	
Indian Spinach young	21	-35	286-417		30	-35
Kang Kong young	16	-28	167-435		15	-32
Tomato						
fruits	16	-37	62-390		9.2-32	

Table 4. Fresh marketable product-nutrient uptake relations of (sub)tropical crops at minimum and maximum N, P and K concentration (kg fresh product per kg nutrient uptake).

crop	N		P		K	
	max-min		max-min		max-min	
Banana/Plantains						
fruits	114	-246	943-2224		24	- 96
Cabbage (white cabbage)						
heads + core's	197	-407	1214-3957		159	-577
Carrot						
roots	174	-853	850-5100		127	-681
Castor (dwarf varieties)						
seeds	8.5-	20	32-	39	9.6-	34
oil	3.6-	8.5	13-	61	4.0-	14
Chillies						
pods (red ripe)	12	- 39	95-	280	9.8-	33
Garlic						
bulbs	34	- 90	278-	638	32	- 96
Okra						
pods	84	-306	617-2231		59	-251
Pigeon Pea						
seeds	7.4-	17	64-	208	15	- 46
Pineapple						
fruits	65	-325	1270-3314		60	-335
Spinach: edible leaves + stems *						
Amaranth	125	-288	863-1150		89	-192
flowering Amaranth	26	-143	167- 526		19	- 28
Indian Spinach	297	-510	4086-5957		433	-493
Kang Kong	177	-309	1856-4833		169	-227
Tomato						
fruits	259	-623	1036-6495		153	-538

* young plants unless indicated otherwise

Appendix 1. References table 1 and 2.

Literature Banana and Plantains

- Awan, J.A. (1978) Horticultural Abstracts 50: 8506.
- Baillon, A.F. et al (1933) Tropical Agriculture Trin. 10: 139-146.
- Bruennich, J.C. (1911/12) on cit by A. Jacob et al (1958) Fertilizer use. Hannover, 617 pp.
- Hey et al (1960) The soils under shifting cultivation.
- Irizarry, H. et al (1981) Journal of Agriculture, University of Puerto Rico 65: 331-345.
- Jauhari, O.S. et al (1974) Indian Journal of Agricultural Chemistry 7: 73-79.
- Loescke, H.W. von (1949) Bananas: chemistry, physiology and technology, pp 20-23.
- Marchal, J. et al (1979) Fruits 34: 373-392.
- Martin-Prevel, P et al (1968) Fruits 23: 259-269.
- Martin-Prevel, P et al (1966) Fruits 21: 19-36.
- Martin-Prevel, P. (1964) Fertilite 22: 3-14.
- Martin-Prevel, P et al (1962) Fruits 17: 123-128.
- Samuels, G. et al (1978) Journal of Agriculture, University of Puerto Rico 62: 178-185.
- Turner, D.W. et al (1985) Scientia Horticulturae 26: 311-322.
- Turner, D.W. et al (1983) Fertilizer Research 4: 89-99.
- Turner, D.W. et al (1982) Scientia Horticulturae 16: 239-252
- Turner, D.W. et al (1980) Scientia Horticulturae 12: 27-45.
- Twyford, I.T. (1973) Plant and Soil 39: 227-243.

Literature White Cabbage

- Abdulla, I. et al (1963) Journal of Agricultural and Food Chemistry 14: 98-109.
- Babish J.G. et al (1979) Journal of Agricultural and Food Chemistry 27: 399-402.
- Beeson, K.C. (1941) on cit by H.D. Chapman (1965) Diagnostic Criteria for Plants and Soils.

- Furr, A.K. (1976) Journal of Food and Agricultural Chemistry 24: 885-888.
- Geissler, T. et al (1971) Archiv fuer Gartenbau 19: 425-441.
- Geissler, T. et al (1953) on cit by Nieuwhof (1969).
- Hara, T. et al (1981) Journal of the Japanese Society of Horticultural Science 50: 60-65.
- Howard, F.D. et al (1962) Californian Agricultural Experiment Station Bulletin no. 788.
- Janes, B.E. (1951) on cit by Nieuwhof (1969).
- Lorenz, O.A. et al (1980) in F.E. Khasawneh et al (eds) The role of Phosphorus in Agriculture pp 737-761.
- Lukovinikova, G.A. et al (1966) Horticultural Abstracts 36: 826.
- Maass, G. et al (1976) Archiv fuer Gartenbau 24: 375-386.
- Nieuwhof, M. (1969) Cole Crops: botany, cultivation and utilization. London.
- Peck, N.H. (1981) Agronomy Journal 65: 679-684.
- Peck, N.H. (1978) Journal of the American Society of Horticultural Science 102: 809-812.
- Peck, N.H. et al (1970) N.Y. State Agric. Exp. Station. Bull. 830.
- Pierre, W.H. et al (1973) Agronomy Journal 65: 91-96.
- Remy, T. (1916) on cit by Nieuwhof (1969).
- Rennie, D.A. (1975) Canadian Journal of Plant Science 55: 761-769.
- Schuphan, W. et al (1985) Qualitas Plantarum et Materiae Vegetabilis 12: 15-47.
- Vogel, F. et al (1938) on cit by Nieuwhof (1969).
- Welch, N.C. et al (1985) Californian Agriculture 39 (7/8): 30-31.

Literature Carrot

- Anonymous (1986) Gemusse 22: 2, 58-61.
- Beeson, K.C. (1941) on cit by Chapman (1965)
- Bernstein, L. et al (1974) Agronomy Journal 66: 412-421.
- Bishop, R.F. et al (1973) Communications in Soil Science and Plant Analysis 4: 455-474.

- Chapman, H.D. (1965) Diagnostic criteria for plants and soils.
- Furr, A.K. (1976) Journal of Agricultural and Food Chemistry 24: 885-888.
- Geyer, B. (1978) Archiv fuer die Gartenbau Berlin 26: 1-13.
- Greenwood, D.J. et al (1980) Journal Agricultural Science Cambridge 95: 457-469.
- Haag, H.P. (1973) Horticultural Abstracts 43: 6109.
- Hamilton, H.A. et al (1975) Canadian Journal of Plant Science 55: 453-461.
- Harrison, H.C. (1986) Journal of the American Society of Horticultural Science 111: 211-215.
- Haworth, F. (1966) Journal Horticultural Science 41: 299-310.
- Hipp, B.W. (1970) HortScience 13: 43-44.
- Howard, F.D. et al (1962) Californian Agricultural Experiment Station Bulletin no 788.
- Howlett, F.S. (1961) on cit by Chapman (1965)
- ILACO (1981) Agricultural compendium for rural development in the tropics and subtropics. Amsterdam, 739 pp.
- Krarup, H. (1985) Horticultural Abstracts 55: 2719.
- Kumar, J.C. et al (1974) Indian Journal of Horticulture 31: 262-267.
- Leh, H.O. (1970) Phytopathologisch Zeitschrift 67: 183-186.
- Linser, H. et al (1975) Zeitschrift fuer Pflanzenernahrung und Bodenkunde 137: 181-196.
- Lorenz, O.A. et al (1980) in F.E. Khasawneh et al (eds) The role of Phosphorus in Agriculture pp 737-761.
- Mauer, A.R. (1971) Canadian Journal of Plant Science 51: 275-281.
- Maynard, D.M. et al (1961) Proceedings of the American Society of Horticultural Science 70: 339-342.
- Nicolaisen, W von et al (1964) Gartenbauwissenschaft 29: 463-480.
- Pankov, V.V. (1978) Horticultural Abstracts 48: 2471.
- Pankov, V.V. (1977) Horticultural Abstracts 47: 5692.
- Rennie, D.A. et al (1975) Canadian Journal of Plant Science 55: 761-769.
- Schnetzer, H.L. (1984) Schweizerische Landwirtschaftliche Forschung 23: 303-314.

Southards, C.J. et al (1962) Proceedings of the American Society of Horticultural Science 81: 335-340.

Vereecken, M. et al (1979) Acta Horticulturae 93: 197-208.

Wunsch, A. (1975) Landwirtschaftliche Forschung 28: 345-358.

Zeid, F.A. et al (1973) Zeitschrift fuer Pflanzenernahrung und Bodenkunde 135: 226-239.

Literature Castor

Allen, S. et al (1985) Journal of Experimental Botany 36: 413-425.

Daussant, J. (1976) Journal of Agricultural and Food Chemistry 24: 103-107.

Hocking, P.J. (1982) Annals of Botany 49: 51-62.

Moshkin, V.A. (1986) Castor. Rotterdam, 315 pp.

Nakagawa, J. et al (1973) Revista de Agricultura 28: 21-30.

Nakagawa, J. et al (1971) Anais da Escola Superior de Agricole "Luiz de Queiroz" 28: 323-337.

Weiss, E.A. (1971) Oil seed crops. London, 901 pp.

Youssef, A.M. et al (1970) Bulletin of the Faculty of Agriculture 18: 253-261.

Zimmermann, L.H. (1958) Advances in Agronomy 10: 257-280.

Literature Chillies

Albasel, N. et al (1984) Journal of Plant Nutrition 7: 1005-1018.

Bagyaraj, D.J. et al (1982) Plant and Soil 69: 375-381.

Beese, F. et al (1982) Agronomy Journal 74: 556-561.

Dadhich, K.S. et al (1969) Plant and Soil 31: 377-379.

El Sherbim, A.H. (1980) Response of paprika to different K levels. PhD-thesis.

Ferrari, J.P. (1972) Journal d' Agriculture tropical et du Botanique applique 19: 63-67.

Govindarajan, V.S. (1985) Critical Reviews in Food Science and Nutrition 22: 109-176.

- Hoskins, M.D. (1976) New Mexico State University Agricultural Experiment Station Research Report 329.
- Howard, F.D. et al (1962) Californian Agricultural Experiment Station Bulletin 788.
- Kariman-Terani, D. et al (1983) *Acta Alimentaria* 12: 301-318.
- Kaufmann, H.G. et al (1971) *Archiv Gartenbau* 19: 6-27.
- Knott, J.E. et al (1967) Vegetable Production in S.E. Asia. Los Banos, 366 pp.
- Lewis, Y.S. (1984) Spices and Herbs for the Food Industry. Orpington, 208 pp.
- Maistre, J. (1964) Les Plantes a epices. Paris, 289 pp.
- Maurya, K.R. et al (1984) *Anales de Edafologia y Agrobiologia* 43: 827-834.
- Miller, C.H. et al (1979) *Journal American Society of Horticultural Science* 104: 852-857.
- Miller, C.H. et al (1961) *Proceedings American Journal of Horticultural Science* 77: 440-448.
- Misra, A. et al (1972) *Indian Journal of Plant Physiology* 15: 56-58.
- Sarudi, J. et al (1939) on cit by Somos (1984).
- Somos, A. (1984) The Paprika.
- Spaldon, E. et al (1968) *Kali Briefe* 16: 40.
- Thomas, J.R. et al (1967) *Agronomy Journal* 59: 27-30.
- Viragh, K. (1981) *Acta Botanica Academicae Scientiarum Hongaricae* 27: 223-264.
- Westphal, E. (1976) Spices, Universite de Yaonde, 52 pp.
- Yamaguchi, M. (1983) World Vegetables, Chichister, 415 pp.

Literature Garlic

- Buwalda, J.G. (1986) *Scientia Horticulturae* 29: 55-68.
- Fenwick, G.F. et al (1985) *Critical Reviews in Food Science and Nutrition* 22: 273-377.
- Lorenz, O.A. et al (1980) in F.E. Khasawneh et al (eds) *The role of Phosphorus in Agriculture* pp 737-761.
- Minard, H.R.G. (1978) *New Zealand Journal of Experimental Agriculture* 6: 139-143.

- Omar, F.A. (1979) Agricultural Research Review 57 (3) 233-241.
Pellet, P.L. et al (1970) on cit by Fenwick et al (1985).
Ramirez, H.V.E. et al (1975) Horticultural Abstracts 45: 7294.
Ruiz, S.R. (1985) Horticultural Abstracts 55: 9460.
USDA (1976) Agricultural Statistics. Washington.
Watt, B.K. (1963) on cit by Fenwick et al (1985).
Zink, F.W. (1963) Journal of the American Society of
Horticultural Science 83: 579-584.

Literature Okra

- Ahmad, N. et al (1968) Agronomy Journal 60: 353-356.
Asif, M.I. et al (1972) Journal of the American Society of
Horticultural Science 97: 440-442.
Costa, M.C.B. et al (1972) Anais da Escola Superior de Agricole
'Luiz de Queiroz' 29: 109-125.
Hipp, B.W. et al (1972) Field Crop Abstracts 42: 7854.
Oyolu, C. (1983) Acta Horticulturae 123: 105-115.
Paliwal, K.V. et al (1972) Journal of Horticultural Science 47:
517-524.
Sharma, J.P. et al (1973) Gartenbauwissenschaft 38: 163-169.
Siemonsma, J.S. (1982) La Culture du Gambo. Diss. Agricultural
University Wageningen, 297 pp.
Singh, R. et al (1974) Current Science 43: 89-91.

Literature Pigeonpea

- Abrams, R. (1975) International Workshop on Grain Legumes, pp
141-147.
Ahlawat, I.P.S. et al (1983) Zeitschrift fuer Acker- und
Pflanzenbau 152: 270-278.
Ahlawat, I.P.S. et al (1981) Journal of Agricultural Research
Camb. 97: 119-124.
Bressani, R. et al (1980) in R.J. Summerfield et al (eds)
Advances in Legume Science.

- Dahiya, B.S. (1977) Field Crop Abstracts 30: 2817.
- Dahiya, B.S. (1977) Field Crop Abstracts 30: 5437.
- Dahiya, B.S. (1977) Qualitas Plantarum, Plant Food for Human Nutrition 27: 327-334.
- Dalal, R.C. (1980) Tropical Agriculture (Trin.) 57: 107-112.
- Dalal, R.C. (1974) Experimental Agriculture 10: 219-224.
- Devarajan, R. et al (1980) Indian Journal of Agricultural Research 14: 47-52.
- FAO (1977) Plant Production and Protection Paper no 9.
- Hedge, D.M. et al (1982) Zeitschrift Acker und Pflanzenbau 151: 302-314.
- ICRISAT (1975) Annual Report for 1975/1976, p 98.
- Irizarry, H. et al (1983) Journal of Agriculture, University of Puerto Rico 67: 188-196.
- Jambunathan, B. et al (1982) in H.C. Srivastava et al (eds) Pulse Production: constraints and prospects, pp 389-395.
- Kumar, M.N.A. et al (1980) Field Crop Abstracts 33: 1426.
- Manimekalai, G. et al (1979) Madras Agricultural Journal 66: 812-816.
- Ramanathan, G. et al (1977) Madras Agricultural Journal 64: 671-672.
- Rao, A.S. et al (1984) Journal of Experimental Botany 35: 774-784.
- Rao, K.J.V.D.K. (1983) Experimental Agriculture 19: 131-141.
- Rao, M.R. et al (1981) Field Crop Abstracts 34: 4626.
- Rathaswamy, R. et al (1973) Madras Agricultural Journal 60: 396-398.
- Salunkhe, D.K. et al (1986) Critical Reviews in Food Science and Nutrition 23: 103-145.
- Sheldrake, A.R. et al (1979) Journal of Agricultural Research Camb. 92: 513-526.
- Singh, B. et al (1980) Legume Research 3: 102-106.
- Singh, S. et al (1968) Cereal Chemistry 45: 13-18.
- Singh, V. et al (1982) Qualitas Plantarum, Plant Food for Human Nutrition 31: 347-354.
- Sinha, S.K. (1982) Field Crop Abstracts 35: 9266.

- Tripathi, A. et al (1982) Field Crop Abstracts 35: 9062.
Tripathi, A. et al (1981) Field Crop Abstracts 34: 8370.
Tripathi, R.D. et al (1977) Field Crop Abstracts 30: 7712.
Wilson, J.R. et al (1983) Australian Journal of the Institute of Agricultural Science 49: 167-169.
Yadav, S.P. (1983) Journal of Agricultural and Food Chemistry 31: 1360-1362.

Literature Pineapple

- Black, R.F. et al (1969) Queensland Journal of Agricultural and Animal Science 26: 385-405.
Boland, F.E. et al (1979) Journal of the Association of Official Agricultural Chemists 55: 200-201.
Dull, G.D. (1971) in A.C. Hulme (ed) The biochemistry of fruits and their products. London, 787 pp.
Lacoeuilhe, J.J. (1979) Kali Briefe 27: 89.
Lacoeuilhe, J.J. (1974) Proceedings of the 10th colloquium of the International Potassium Institute 1973: 177-183.
Lacoeuilhe, J.J. et al (1971) Fruits 26: 581-597.
Lodh, S.B. (1972) Indian Journal of Horticulture 29: 287-291.
Norman, J.C. (1980) Beitrage zuer Tropischen Landwirtschaft und Veterinaermedizin 18: 333-334.
Onaha, A. et al (1986) Journal of the Japanese Society of Horticultural Science 54: 438-449.
Page, P.E. (1971) Queensland Journal of Agricultural and Animal Science 28: 195-197.
Page, P.E. (1971) Queensland Journal of Agricultural and Animal Science 28: 192-194.
Sale, J.W. (1938) Journal of the Association of Official Agricultural Chemists 21: 502-515.
Sideris, C.P. et al (1946) Plant Physiology 21: 247-270.
Sideris, C.P. et al (1945) Plant Physiology 20: 609-639.
Teisson, C. et al (1982) Fruits 37: 741-748.

Literature Spinach

- Cornelis, J. et al (1985) Abstracts on Tropical Agriculture 10 (4) 9-21.
- Donald, E.L. (1986) Horticultural Abstracts 56: 6164.
- Faboya, O.O.P. (1983) Food Chemistry 12: 213-216.
- Grubben, G.J.M. (1975) Mededelingen Landbouw Hogeschool Wageningen 75: 6.
- Hill, R.M. et al (1982) Journal of Agricultural and Food Chemistry 30: 465-469.
- Ifon, E.T. et al (1979) Food Chemistry 4: 263-267.
- Johnson, C.R. (1973) Communications in Soil Science and Plant Analysis 4: 185-196.
- Katiyar, S.K. et al (1985) Journal of Food Science and Technology 22: 438-440.
- Linnemann, A.R. et al (1986) Netherlands Journal of Agricultural Science 34: 503-505.
- Norman, J.C. (1972) World Crops 24: 217-219.
- Oliveira, J.S. et al (1975) Economic Botany 29: 255-263.
- Olufolaji, A.O. et al (1980) Scientia Horticulturae 13: 181-189.
- Purushothaman, V. et al (1980) Vegetables for the hot, humid Tropics 5: 26-38.
- Schmidt, D.R. (1971) Agronomy Journal 63: 546-550.
- Sheeramulu, N. (1982) Journal of Plant Foods 4: 139-141.
- Stafford, W.I. (1976) Plant Foods for Man 2: 7-13.
- Taylor, O.A. (1982/83) Scientia Horticulturae 18: 313-322.
- Vuurmans, J. et al (1977) Vegetables for the hot, humid Tropics 2: 40-43.
- Wills, R.B.H. et al (1984) Journal of Agricultural and Food Chemistry 32: 413-416.

Literature Tomato

- Aung, L.H. (1983) Journal of Agricultural and Food Chemistry 31: 1259-1262.
- Beeson, K.C. (1941) on cit by H.D. Chapman (1965) Diagnostic criteria for plants and soils.

- Besford, R.T. (1979) Plant and Soil 51: 331-340.
- Besford, R.T. (1978) Plant and Soil 50: 179-191.
- Besford, R.T. et al (1975) Plant and Soil 42: 395-412.
- Cooper, A.J. (1955) Proceedings 14th International Horticultural Congress, Vol 1: 589-595.
- Erickson, A.B. et al (1982) Plant growth and regulation 1: 11-14.
- Furr, A.K. (1976) Journal of Agricultural and Food Chemistry 24: 885-888.
- Graifenberg, A. et al (1986) Acta Horticulturae 176: 83-91.
- Granges, A. (1980) Revue Suisse de Viticulture, d'Arboriculture et de Horticulture 12: 59-63.
- Greidanus, P. et al (1971) Mededelingen van het Sprenger Instituut, no 30.
- Hartman, P.L. (1986) Journal of the American Society of Horticultural Science 103: 265-269.
- Hermann, K. (1979) Zeitschrift Lebensmitteluntersuchung und Forschung 169: 179-200.
- Hills, F.J. (1983) Agronomy Journal 75: 423-426.
- Ho, L.C. (1984) Plant growth and regulation 2: 277-285.
- Hurd, R.G. et al (1979) Annals of Applied Biology 93: 77-89.
- Kaufmann, H.G. et al (1968) Archiv fuer die Gartenbau 16: 359-377.
- Kidson, E.B. et al (1953) New Zealand Journal of Science and Technology 35: 127.
- Kirby, E.A. (1967) Plant Physiology 42: 6-14.
- Lorenz, O.A. et al (1980) in F.E. Khasawneh et al (eds) The role of Phosphorus in Agriculture pp 737-761.
- Maher, M.J. (1976) Scientia Horticulturae 4: 23-26.
- Mayr, H.H. (1969) in K. Scharrer et al (eds) Handbuch der Pflanzenernahrung und Dunung, pp 9-89.
- Orphanos, P.I. et al (1980) Journal Horticultural Science 55: 415-423.
- O'Sullivan, J. et al (1974) Journal of the American Society of Horticultural Science 99: 543-547.
- Oswiecimski, I. (1982) Horticultural Abstracts 52: 1572.
- Pill, W.G. et al (1980) Journal of the American Society of Horticultural Science 105: 730-734.

- Pill, W.G. et al (1979) Canadian Journal of Plant Science 59: 391-397.
- Pill, W.G. et al (1978) Journal of the American Society of Horticultural Science 103: 265-269.
- Roorda, J.P.N.L. et al (1981) Nutritional disorders in glasshouse tomatoes, cucumber and lettuce. Wageningen, 130 pp.
- Souci, S.W. (1962) on cit by Hermann (1979).
- Tanaka, A. et al (1974) Soil Science and Plant Nutrition 20: 57-68.
- Uexhull, H.R. et al (1978) First International Symposium on Tropical Tomato. AVRDC publication no 78-59, pp 65-78.
- Ward, G.M. (1967) Proceedings of the American Society of Horticultural Science 90: 335-341.
- Watt, B.K. (1963) on cit by Hermann (1979).
- Widders, I.E. et al (1982) Journal of the American Society of Horticultural Science 107: 960-964.
- Widders, I.E. et al (1979) Journal of the American Society of Horticultural Science 104: 216-220.

Appendix 2.

Minimum and maximum N, P and K concentrations in economic products and crop residues of (sub)tropical crops obtained from the literature. Concentrations in the residues are between brackets if the concentrations in the residues and economic products are from different crops. Harvest Indices (HI) are expressed in dry matter and in percentages. Haulms are defined as stems and leaves. Shelling is the seed/pod percentage. D.M. % indicates the dry matter concentration.

Reference	N min-max %	P min-max %	K min-max %
Crop crop part			
Banana and Plantains			
Awan (1978)			
fruits	0.91-0.91		
Baillon et al (1933)			
fruits	1.02-1.02	0.12-0.12	2.96-2.96
Bruennich (1911/12)			
bunch	0.88-1.00	0.12-0.14	2.53-3.49
residue	0.87-1.33	0.08-0.12	2.53-2.53
HI (bunch)	39.9		
Hye et al (1970)			
DM % fruits	30		
Irizarry et al (1981)			
bunch	0.66-0.66	0.08-0.08	1.72-1.72
residue	0.96-0.96	0.06-0.06	2.04-2.04
HI (bunch)	44.6		
Jauhari et al (1974)			
residue	2.78-?..?	0.55-?..?	6.72-?..?
Loescke (1949)			
DM % fruits	20.1		
Marchal et al (1979)			
bunch	0.60-0.87	0.09-0.12	2.75-2.76
fruits	0.58-0.85	0.09-0.11	2.51-2.51
residue	0.74-1.18	0.07-0.11	4.74-8.96
HI (bunch)	51.5		
HI (fruits)	50.1		
Martin-Prevel et al (1968)			
bunch	1.26-1.26	0.18-0.18	3.65-3.65
fruits	1.26-1.26	0.17-0.17	3.31-3.31
residue	1.00-1.00	0.16-0.16	5.03-5.03
HI (bunch)	28.6		
HI (fruits)	27.7		
HI (fruits)	22.9 (incl shed leaves)		
DM % fruits	16.6 (approximation)		

Reference	N min-max %	P min-max %	K min-max %
Crop crop part			
Martin-Prevel et al (1966)			
fruits	0.79-1.18	0.09-0.14	1.90-2.54
Martin-Prevel (1964)			
bunch	0.59-1.33	0.11-0.14	2.91-4.11
fruits	0.59-1.27	0.10-0.13	2.25-3.31
fruit wgt/bunch weight	0.953		
Martin-Prevel et al (1962)			
bunch	1.11-1.11	0.16-0.16	4.12-4.12
residue	1.37-1.37	0.10-0.10	6.22-6.22
HI (bunch)	46.9		
HI (bunch)	44.2 (incl shed leaves)		
DM % fruits	18.5		
Samuels et al (1978)			
fruits	0.64-0.64	0.08-0.08	2.22-2.22
residue	1.44-1.44	0.07-0.07	8.09-8.09
HI (fruits)	37.5		
Turner et al (1985)			
residue	1.76-?..?	0.22-?..?	3.40-?..?
Turner et al (1983) approximation			
bunch	1.33-1.33	0.18-0.18	2.58-2.58
fruits	1.49-1.49	0.16-0.16	2.79-2.79
residue	1.03-1.03	0.17-0.17	1.44-1.44
Turner et al (1982)			
fruits	0.89-1.83	0.14-0.18	2.39-3.29
Turner (1980)			
HI (bunch)	28.9		
HI (fruits)	28.3		
Twyford (1973)			
bunch	0.71-1.28	0.09-0.14	2.77-2.77
fruits	0.71-1.25	0.10-0.14	2.62-2.62
residue	0.75-0.88	0.12-0.17	1.57-2.65
HI (bunch)	28.9		
HI (fruits)	28.0		
HI (fruits)	26.7		
DM % fruits	16.3 (approximation)		
White Cabbage			
Abdulla et al (1963)			
straw		0.13-?..?	
Babish et al (1979)			
core + head	2.33-3.54	0.32-0.53	1.84-2.96
Beeson (1941)			
core + heads		0.13-0.77	
Furr (1976)			
core + heads			2.73-3.12
Geissler et al (1971)			
core + heads	2.64-2.82		

Reference	N min-max %	P min-max %	K min-max %
Crop crop part			
<hr/>			
Geissler et al (1953)			
core + heads			3.60-3.70
Hara et al (1981)			
core + heads	2.09-?..??		
straw	1.36-?..??		
HI (heads + core's)	32.7 or more (immature plants)		
Howard et al (1962)			
core + heads	2.40-2.40	0.43-0.43	2.75-2.75
Lorenz (1980)			
core + head		0.38-0.38	
Janes (1951)			
core + heads		0.50-0.70	2.20-3.70
Lukovinikova et al (1966)			
core + heads	2.52-2.52		
Maass et al (1976)			
core + heads	2.76-3.10	0.42-0.47	3.28-3.61
Nieuwhof (1969)			
core + heads	3.40-3.40	0.50-0.50	3.90-3.90
straw (approximation)	1.87-1.87	0.38-0.38	3.78-3.78
HI	47.9 (author's interpretation)		
DM % heads	7.1		
Peck (1981)			
core + heads	1.26-1.82		
straw	1.38-2.59		
HI	58.6		
HI (incl. shed leaves)	57.6 or less		
Peck (1978)			
core + heads (approximation)		0.31-0.44	1.60-4.05
straw		0.19-0.19	1.18-1.18
HI (approximation)	47.6		
Peck (1970)			
core + head		0.23-0.38	1.31-3.55
straw		0.17-0.47	0.66-3.43
HI	61.6		
HI (incl. shed leaves)	61.1		
DM % heads	6.4		
Pierre et al (1973)			
core + heads	2.86-3.93		
Remy (1916)			
core + heads	2.90-2.90	0.50-0.50	3.30-3.30
Rennie (1975)			
core + heads	2.22-2.82	0.29-0.36	
HI	33.4		
Schuphan et al (1985)			
core + heads	4.20-4.60	0.45-0.49	2.35-2.38
Vogel et al (1938)			
core + heads	2.90-3.70	0.30-0.50	4.10-5.10
Welch et al (1985)			
straw	1.89-?..??	0.52-?..??	3.25-?..??

Reference	N min-max %	P min-max %	K min-max %
Crop crop part			
Carrot			
Anonymous (1986)			
root	0.67-1.12	0.27-0.34	2.37-2.68
Beeson (1941)			
root		0.14-0.65	
Bernstein et al (1974)			
top	2.00-?..?	0.20-?..?	1.88-?..?
Bishop et al (1973)			
top	1.82-3.54	0.16-0.60	1.02-6.20
Furr (1976)			
root			2.53-2.94
Geyer (1978)			
root	1.85-1.85		
Greenwood et al (1980)			
root		0.32-0.36	
top		0.15-0.18	
HI	73.0		
Haag (1973)			
top	0.79-?..?	0.09-?..?	
Hamilton et al (1975)			
root	2.14-2.40	0.49-0.49	2.52-2.84
top	2.54-2.62	0.54-0.54	2.68-3.14
HI	57.6		
DM % roots	10.6		
Harrison (1986)			
root	1.21-1.81	0.27-0.40	1.56-2.85
Haworth (1966)			
root	1.30-1.82	0.20-0.29	0.93-3.22
top	2.15-2.82	0.16-0.34	1.12-4.32
Hipp (1970)			
top	2.46-?..?		
Howard et al (1962)			
root	1.16-1.16	0.33-0.33	2.27-2.27
Howlett (1961)			
top		0.26-0.26	
ILACO (1981)			
DM % roots	11		
Krarup (1985)			
root	0.65-1.02	0.06-0.06	0.56-0.69
top	(1.68-1.99)	(0.05-0.05)	(0.45-0.63)
Kumar et al (1974)			
root	0.88-1.22		
DM % root	9.6		
Leh (1970)			
root	2.00-2.00		3.87-4.38
top	1.81-2.74		2.13-5.68

Reference	N min-max %	P min-max %	K min-max %
Crop crop part			
<hr/>			
Linser et al (1975)			
root	0.48-0.58		
top	0.52-0.60		
HI	72.8		
HI (incl. fallen leaves)	52.8		
Lorenz (1980)			
root		0.33-0.33	
top		(0.20-0.35)	
Mauer (1971)			
root	1.46-1.79	0.34-0.38	2.85-3.39
top	2.41-2.68	0.37-0.42	3.86-4.75
Maynard et al (1961)			
root			3.91-4.21
Nicolaisen, von et al (1964)			
root	0.86-1.81		
top	1.75-3.91		
HI	78.1		
Pankov (1978)			
top			1.70-?..??
Pankov (1977)			
top		0.25-?..??	
Rennie et al (1975)			
root	1.33-2.00	0.20-0.33	
top	2.54-2.58	0.25-0.26	
HI	60.6		
Schnetzer (1984)			
root	0.42-1.49		
top	1.41-2.39		
Southards et al (1962)			
top	1.39-3.95	0.18-0.66	0.98-6.52
Vereecken et al (1979)			
top	4.06-?..??	0.28-?..??	
Wunsch (1975)			
root	0.91-1.42		
top	2.27-2.54		
Zeid et al (1973)			
root	0.58-0.58		
top	0.68-0.68		
HI	76.8		
<hr/>			
Castor (dwarf varieties)			
Allen et al (1985)			
straw	5.33-?..??	1.48-?..??	10.30-?..??
Daussant (1976)			
seeds	2.88-3.20		

Reference Crop crop part	N min-max %	P min-max %	K min-max %
Hocking (1982)			
leaves	2.98-4.56	0.22-0.31	1.65-2.05
seeds	2.88-2.88	0.60-0.60	0.57-0.57
pods	2.10-2.10	0.35-0.35	2.42-2.42
shelling %	48.6		
DM % seeds	85.1		
Moshkin (1986)			
seeds	3.00-3.00		
pods approximation	2.07-2.07		
straw	2.36-?.	0.18-?.	1.93-?.
HI (pods)	26		
HI (seeds)	21		
Nakagawa et al (1973)			
pods	2.39-2.39	0.28-0.28	0.43-0.43
straw	2.80-?.	0.09-?.	2.57-?.
Nakagawa et al (1971)			
straw	2.28-?.	0.07-?.	2.45-?.
Weiss (1971)			
shelling %	61		
oil % seeds	48		
Youssef et al (1970)			
seeds	3.98-4.28	0.79-0.83	
Zimmermann (1958)			
shelling %	65		
oil % seeds	50		
Chillies			
Bagyaraj et al (1982)			
straw		0.11-0.20	
HI (rough approximation)	34.7		
Beese et al (1982)			
HI	61.9		
Dadhich et al (1969)			
straw	2.00-3.03		
HI	32.1		
Ferrari (1972)			
fruits	2.90-3.33		
Govindarajan (1983)			
fruits	1.91-2.83	0.39-0.39	
DM % pods	90		
Hoskins (1976)			
fruits			2.73-4.02
Howard et al (1962)			
fruits	2.29-2.29	0.33-0.33	1.86-1.86
Kariman-Terani et al (1983)			
fruits			1.88-1.88
straw			1.59-2.08

Reference Crop crop part	N min-max %	P min-max %	K min-max %
Knott et al (1967) fruits	3.19-3.68	0.27-0.50	1.25-1.42
Lewis (1984) fruits	2.58-2.58		
Maistre (1964) fruits	2.75-2.75	0.46-0.46	3.01-3.01
Maurya et al (1984) fruits	3.19-3.68	0.24-0.27	1.25-1.42
Misra et al (1972) fruits	1.49-1.54		
Sarudi et al (1939) fruits		0.41-0.41	3.39-3.39
Viragh (1980) HI	62.5		
Westphal (1976) fruits	2.82-2.82		
Yamaguchi (1983) fruits	2.00-2.29	0.28-0.33	
Other Capsicum spp. (sweet and bell peppers)			
Albasel et al (1984) fruits	1.90-2.70		
straw	1.38-1.85		
El Sherbim (1980) fruits	3.34-3.34	0.46-0.46	3.46-3.46
straw	2.93-2.93	0.33-0.33	5.91-5.91
Kaufmann et al (1971) fruits	2.82-3.01	0.37-0.39	3.18-3.38
straw	(2.54-2.71)	(0.18-0.21)	(2.94-3.12)
Miller et al (1979) fruits	2.20-2.20	0.32-0.32	2.50-2.50
straw	2.30-2.30	0.35-0.35	2.80-2.80
Miller et al (1961) fruits	1.58-3.30	0.22-0.44	1.84-3.38
straw	1.09-4.93	0.12-0.40	1.00-4.64
Somos (1984) fruits	2.39-3.00	0.40-0.61	1.91-3.19
straw	(1.32-3.85)	(0.22-0.65)	(2.04-5.29)
Spaldon et al (1968) fruits	4.06-4.74	0.21-0.76	5.12-6.95
straw	3.13-3.19	0.11-0.32	3.95-3.98
Thomas et al (1967) fruits	2.25-2.47	0.34-0.46	
straw	2.24-2.24	0.18-0.18	

Reference	N min-max %	P min-max %	K min-max %
Crop crop part			
Garlic			
Buwalda (1986)			
bulb	0.05-0.81		
top	0.26-1.20		
HI	52.1		
Fenwick (1985)			
DM % bulbs	37.5		
Lorenz (1980)			
bulbs		0.38-0.38	
Minard (1978)			
bulb	1.90-2.30	0.26-0.34	1.00-1.20
top	3.05-3.55	0.26-0.32	3.70-4.70
DM % bulbs	32		
Omar (1979)			
bulb	1.00-1.15		
top	2.08-3.10		
Pellet et al (1970)			
bulb	2.34-2.34		
Ramirez et al (1975)			
top		0.40-?.??	
Ruiz (1985)			
top	5.50-?.??	0.50-?.??	2.00-?.??
USDA (1976)			
bulb		0.38-0.38	
Watt (1963)			
bulb	2.56-2.56	0.52-0.52	1.37-1.37
Zink (1963)			
bulb	2.14-2.66	0.31-0.41	0.89-1.03
top	(1.95-?.???)	(0.28-?.???)	(2.20-?.???)
DM % bulbs	41.8		
Okra			
Ahmad et al (1968)			
pods	2.10-3.49	0.53-0.71	2.10-2.80
Asif et al (1972)			
pods	2.19-2.19	0.49-0.49	1.94-2.10
Costa et al (1972)			
pods	3.15-3.15	0.54-0.55	1.90-4.40
straw	1.96-2.45	0.22-0.27	1.53-3.86
HI (pods)	28.9		
Hipp et al (1972)			
straw		0.30-?.??	
Oyolu (1983)			
pods	2.22-2.22	0.40-0.40	0.28-0.28
Paliwal et al (1972)			
pods	1.16-2.16	0.26-0.46	1.80-4.18
straw	0.97-?.??	0.05-?.??	1.36-?.??

Reference	N min-max %	P min-max %	K min-max %
Crop crop part			
Sharma et al (1973)			
pods	2.23-2.77		
straw	0.38-? .??		
Siemonsma (1982)			
pods	2.93-3.04	0.38-0.40	2.35-2.63
straw	1.66-1.68	0.16-0.16	0.85-0.87
HI (pods)	22.2		
HI (pods)	19.2 (incl leaves shed)		
DM % pods	10		
Singh et al (1974)			
pods	2.27-2.27	0.17-0.17	
Pigeonpea			
Abrams (1975)			
seeds	3.30-3.49		
Ahlawat et al (1983)			
seeds	3.43-3.58	0.47-0.51	
straw	1.01-1.01	0.12-0.13	
HI (seeds)	27.4		
Ahlawat et al (1981)			
seeds	3.45-3.58	0.43-0.53	
straw	1.01-1.02	0.11-0.15	
HI (seeds)	24.6		
Bressani et al (1980)			
seeds	3.66-3.66		
Dahiya (1977)			
seeds	2.74-3.84		
Dahiya (1977)			
seeds	2.93-4.32		
Dahiya (1977)			
seeds	2.94-4.06		
Dalal (1980)			
seeds	3.17-3.17	0.33-0.33	1.05-1.05
shells	1.47-1.47	0.10-0.10	0.32-0.32
haulms	2.27-2.27	0.17-0.17	0.51-0.51
straw	1.92-1.92	0.14-0.14	0.42-0.42
HI (seeds)	29.9		
shelling %	49.3		
Dalal (1974)			
seeds	3.59-3.59	0.15-0.15	0.94-0.94
straw	1.61-1.61	0.11-0.11	0.60-0.60
HI (seeds)	36.5		
FAO (1977)			
seeds	2.96-4.21		
Hegde et al (1982)			
seeds	3.02-3.27	0.33-0.39	0.63-1.22
shells	0.89-0.89	0.09-0.11	0.35-0.35
haulms	1.10-1.31	0.09-0.14	0.21-0.31
straw	1.06-1.25	0.09-0.13	0.23-0.31

Reference Crop crop part	N min-max %	P min-max %	K min-max %
ICRISAT (1975) haulms	1.40-1.40		
Irizarry et al (1983) pods	3.14-3.14	0.09-0.09	1.68-1.68
haulms	1.38-1.38	0.09-0.09	1.23-1.23
HI (seeds)	18.1		
HI (incl. fallen leaves)	15.9		
shelling %	60.0		
Jambunathan et al (1982) seeds	2.48-4.58		
Kumar et al (1980) shells	1.45-1.45	0.09-0.09	
Manimekalai et al (1979) seeds	3.16-3.91		
Rao et al (1984) seeds	2.96-2.96		
shells	0.78-0.78		
haulms	0.74-0.74		
straw	0.74-0.74		
HI (seeds)	30.0		
HI (incl. fallen leaves)	25.2		
shelling %	68.5		
Rao (1983) seeds	3.03-3.09		
shells	0.67-0.70		
haulms	0.67-0.81		
straw	0.67-0.79		
HI (seeds)	27.9		
shelling %	65.1		
Rao et al (1981) seeds	3.12-3.12	0.50-0.50	
shells	0.78-0.78	0.06-0.06	
Rathaswamy et al (1973) seeds	2.88-3.42		
Salunkhe (1986) seeds	4.06-4.06	0.41-0.41	
DM % seeds	9.85		
Sheldrake et al (1979) seeds	3.38-3.45	0.28-0.29	
shells	0.68-0.68	0.03-0.03	
haulms	0.99-1.13	0.06-0.06	
straw	0.97-1.09	0.05-0.06	
HI (seeds)	22.4		
HI (incl. fallen leaves)	16.0		
shelling %	66.5		
Singh et al (1980) seeds	3.74-4.08		
shelling %	61.1		
Singh et al (1968) seeds	3.66-3.66	0.36-0.36	

Reference		N min-max %	P min-max %	K min-max %
Crop	crop part			
Singh et al (1982)				
seeds		3.28-3.28		
Singha (1982)				
HI (seeds)		30.7		
Tripathi et al (1982)				
seeds		3.18-3.44		
Tripathi et al (1981)				
seeds		2.85-3.41		
Tripathi et al (1977)				
seeds		3.19-4.08		
Wilson et al (1983)				
haulms		3.30-?..??	0.24-?..??	
Yadav (1983)				
seeds		4.42-4.42		
shells		1.14-1.14		
DM % seeds		21.4		
Pineapple				
Black et al (1969)				
fruit		1.12-1.12	0.17-0.17	1.62-1.62
fruit + crown		1.21-1.21	0.16-0.16	1.58-1.58
residue		1.02-1.02	0.09-0.09	1.53-1.53
HI (fruits)		13.2		
HI (fruit + crown)		20.1		
Boland et al (1979)				
fruits		0.39-1.23	0.03-0.13	0.38-2.16
Dull (1971)				
fruits (approximation)		0.28-0.71	0.04-0.16	0.08-2.46
Lacoeuilhe (1979)				
residue		1.42-?..??	0.11-?..??	2.87-?..??
HI		29.2		
Lacoeuilhe (1974)				
residue		1.12-?..??		2.14-?..??
Lacoeuilhe et al (1971)				
fruits		0.52-0.61	0.10-0.11	0.89-1.82
fruit + crown		0.68-0.96	0.12-0.12	0.76-2.07
residue		0.95-1.24	0.14-0.18	0.75-2.98
HI (fruits)		37.2		
HI (fruits + crown)		43.5		
DM % fruits		13.1		
Lodh (1972)				
fruits		0.54-0.54	0.01-0.01	2.08-2.08
DM % fruits		10.6		
Norman (1980)				
HI fruits		9.4		
HI fruit/crown/stalk		32.4		
DM % fruits		4.9		

Reference Crop crop part	N min-max %	P min-max %	K min-max %
Onaha et al (1986)			
HI fruits	24.9		
HI fruit/crown/stalk	29.8		
HI fruits	22.8 incl fallen leaves		
HI fruit/crown/stalk	27.8 incl fallen leaves		
Page (1971)			
residue	0.87-5.30		
Page (1971)			
residue			1.53-4.29
Sale (1938)			
fruits		0.04-0.04	1.04-1.04
Sideris et al (1946)			
residue		0.13-?.??	3.19-?.??
Sideris et al (1945)			
residue		0.07-?.??	1.75-?.??
Teisson et al (1982)			
HI (fruits)	38.5		
HI fruit/crown/stalk	50.4		
DM % fruits	16.5		
Spinach, nutrient concentrations of whole plants			
Cornelis et al (1985) <i>Ipomoea aquatica</i>			
young plant	3.43-6.37	0.04-0.56	4.75-5.21
HI	100		
DM %	9.8		
Grubben (1975) <i>Amaranthus</i> spp.			
young plant	3.51-3.51	0.61-0.61	7.88-7.88
HI	67		
DM % young plant	9.6		
flowering plant	1.63-1.63	0.31-0.31	3.75-3.75
HI	22		
Hill et al (1982) <i>Amaranthus</i> spp.			
flowering plant	3.38-4.29	0.33-0.52	5.43-5.63
Linneman et al (1986) <i>Ipomoea aquatica</i>			3.01-5.68
Olufolaji et al (1980) <i>Amaranthus cruentus</i>			
young plants	2.12-3.45	0.68-0.77	6.05-7.44
flowering plants	0.56-0.93	0.50-0.59	3.58-4.35
HI (approximation)	20		
Purushothaman et al (1980)			
young Kang Kong	4.86-4.86	0.62-0.62	6.76-6.76
HI	100		
DM %	7.6		
Young Amaranth	3.94-3.94	0.68-0.68	5.89-5.89

Reference		N min-max %	P min-max %	K min-max %
Crop	crop part			
Schmidt (1971)				
young Amar. tricolor		4.57-5.28		2.92-4.20
HI (approximation)		50		
DM %		14		
young Basella alba		4.01-4.80		2.92-3.24
HI		100		
DM %		7.1		
young Ipomoea aquatica		5.10-5.70		3.26-4.07
HI		100		
DM %		11.8		
Taylor (1982/83)	Amaranthus caudatus flowering plant		0.16-?..??	4.43-?..??
Vuurmans (1977)				
Indian Spinach young		1.74-3.43		
Kang Kong young		3.08-4.78		
Wills et al (1984)				
young Amaranthus spp.		5.59-5.59		7.59-7.59
HI		89		
DM %		8.3		
DM % Kang Kong		7.8		
Spinach, nutrients in leaves.				
Donald (1986)				
Basella alba		2.63-2.63	0.36-0.36	3.55-3.55
Faboya (1983)				
Amaranthus caudatus			0.30-0.33	0.64-0.77
Basella alba			0.23-0.25	0.50-0.52
Grubben (1975)				
DM %		15.6		
Ifon et al (1979)				
Amaranthus hybridus		0.66-0.68	4.20-4.38	
Johnson (1973)				
Amaranthus spp.				
leaves + stemtops		5.77-5.77	0.61-0.61	8.72-8.72
Katiyar et al (1985)				
Amaranthus polygamus		4.70-4.70	0.72-0.72	
Norman (1972)				
Amaranthus hybridus		5.33-5.33		
Oliveira et al (1975)				
Amaranthus caudatus				
edible leaves		4.28-4.28	0.31-0.31	0.26-0.26
DM % leaves		18.8		
Sheeramuku (1982)				
Amaranthus spp.		3.22-4.59		
Basella alba		4.24-4.24		
Ipomoea aquatica		4.61-4.61		
Stafford (1976)				
Amaranthus hybridus		1.96-5.11		
Taylor (1982/83)				
Amaranthus caudatus			0.20-0.29	4.77-5.35

Reference	N min-max %	P min-max %	K min-max %
Crop crop part			
Tomato			
Aung (1983) fruits		0.06-0.07	1.49-3.03
Beeson (1941) fruits		0.29-0.84	
Besford (1979) fruits		0.17-?.??	
straw		0.05-?.??	
Besford (1978) fruits			0.98-4.83
Besford et al (1975) fruits	2.20-3.42	0.42-0.45	1.50-4.34
Cooper (1955) HI (incl leaves shed)	0.81 * HI (excl leaves shed)		
Author's interpretation			
Erickson (1982) DM % fruits	4.5		
Furr (1976) fruits			2.22-2.51
Graifenberg et al (1986) fruits	1.65-1.65		
straw	1.50-1.50		
HI	62.7		
Granges (1980) fruits	1.77-2.53	0.32-0.50	3.22-4.52
Greidanus et al (1971) fruits	2.71-3.20	0.40-0.42	3.45-6.00
Hartman (1986) fruits	2.61-?.??	0.58-?.??	1.95-?.??
straw	3.03-?.??	0.85-?.??	2.45-?.??
Hills (1983) fruits	2.30-2.30		
straw	1.74-1.74		
HI	62.4		
Ho (1984) DM % fruits	7.7		
Hurd, R.G. (1979) HI	79		
HI (incl shed leaves)	65		
DM % fruits	7		
Kaufmann et al (1968) fruits	2.59-2.71	0.43-0.61	4.38-4.82
straw	2.11-2.42	0.28-0.31	3.27-3.67
HI	44.2		
Kidson et al (1953) fruits	2.17-2.73	0.40-0.54	4.38-6.58
straw	(2.06-2.26)	(0.33-0.55)	(2.58-2.82)
HI	62.5		

Reference		N min-max %	P min-max %	K min-max %
Crop	crop part			
Kirby (1967)				
straw		1.75-?.??		3.60-?.??
Lorenz (1980)			0.55-0.55	
fruits				
Maher (1976)				
fruits		3.00-3.00	0.50-0.50	4.50-4.50
straw		3.12-3.12	0.50-0.50	5.13-5.13
HI (incl shed leaves)		43.0		
Mayr (1969)				
fruits			0.35-0.35	
Orphanos et al (1980)				
fruits		1.83-3.31		
O'Sullivan et al (1974)				
straw		0.74-?.??		
Oswiecimski (1982)				
straw		0.30-?.??	0.17-?.??	2.91-?.??
Pill et al (1980)				
fruits				3.07-4.57
Pill et al (1979)				
fruits			0.54-0.62	4.72-5.35
Pill et al (1978)				
fruits			0.85-0.93	3.02-4.83
Roorda et al (1981)				
fruits		1.68-3.50	0.28-0.80	2.74-5.87
Souci (1962)				
fruits		1.90-2.76	0.30-0.44	4.48-5.34
Tanaka et al (1974)				
fruits		2.78-2.78	0.41-0.41	2.99-2.99
straw		(2.24-2.24)	(0.22-0.22)	(4.01-4.01)
HI		53.6		
HI (incl shed leaves)		48.8		
DM % fruits		6.7		
Uexhull et al (1978)				
fruits		1.93-1.93	0.41-0.41	3.77-3.77
straw		2.34-2.34	0.51-0.51	5.39-5.39
HI		70		
Ward (1967)				
fruits		1.81-1.81	0.42-0.42	3.40-3.40
straw		2.61-2.61	0.80-0.80	3.25-3.25
HI (approximation)		81.0		
Watt (1963)				
fruits		2.71-2.71	0.42-0.42	3.75-3.75
Widders et al (1982)				
fruits				3.67-4.09
straw				2.38-2.38
straw (extreme values)				1.98-2.55
HI		57.0		
Widders et al (1979)				
fruits				3.94-4.09
straw				1.68-3.50
HI		55.5		